w abeomics

36-3585: Anti-CD33 / SIGLEC3 (Myeloid Cell Surface Antigen) Monoclonal Antibody(Clone: WM53)

Clonality :	Monoclonal
Clone Name :	WM53
Application :	FACS,IF,Functional Assay
Reactivity :	Human
Gene :	CD33
Gene ID :	945
Uniprot ID :	P20138
Alternative Name :	gp67; My9; Myeloid cell surface antigen CD33; p67; Sialic acid-binding Ig-like lectin 3 (SIGLEC3)
Isotype :	Mouse lgG1, kappa
Immunogen Information : Human AML cells	

Description

Recognizes a 67kDa glycoprotein, which is identified as CD33 (HLDA IV; WS Code M-505). CD33 is a transmembrane protein of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. It belongs to the immuno-receptor tyrosine-based inhibitory motif (ITIM)-containing molecules able of recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to signal assemblies; these ITIMs are also used for ubiquitin-mediated removal of the receptor from the cell surface. CD33 is expressed on cells of myelomonocytic lineage, binds sialic acid residues in N- and O-glycans on cell surfaces, and is a therapeutic target for acute myeloid leukemia. CD33 is expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. It is absent on platelets, lymphocytes, erythrocytes and hematopoietic stem cells.

Product Info

Amount :	20 μg / 100 μg
Content :	200 μ g/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Functional Studies (Order Ab without BSA & Azide);