

## 36-3590: Anti-CD34 (Hematopoietic Stem Cell & Endothelial Marker) Monoclonal Antibody(Clone: HPCA1/1806R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	HPCA1/1806R
<b>Application :</b>	FACS,IHC
<b>Reactivity :</b>	Human, Rat
<b>Gene :</b>	CD34
<b>Gene ID :</b>	947
<b>Uniprot ID :</b>	P28906
<b>Alternative Name :</b>	Hematopoietic Progenitor Cell Antigen, HPCA1, Mucosialin
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant full-length human HPCA1 protein

### Description

This antibody recognizes a transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi's sarcoma, but shows low specificity.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of recombinant MAb Purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Flow Cytometry (1-2ug/million cells); Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes)




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with CD34 Rabbit Recombinant Monoclonal Antibody (HPCA1/1806R).




Fig. 2: SDS-PAGE Analysis of Purified CD34 Rabbit Recombinant Monoclonal Antibody (HPCA1/1806R).




Fig. 3: Flow Cytometric Analysis of human Jurkat cells using CD34 Rabbit Recombinant Monoclonal Antibody (HPCA1/1806R) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).