

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3621: Anti-CD44v6 (Marker of Tumor Metastasis) Monoclonal Antibody(Clone: CD44v6/1246)

Monoclonal Clonality: CD44v6/1246 Clone Name:

Application: IHC Reactivity: Human Gene: CD44 Gene ID: 960 **Uniprot ID:** P16070

LHR; BA-1; chondroitin sulfate proteoglycan 8 (CSPG8); Epican; Extracellular Matrix Receptor III

(ECM III); GP90 Lymphocyte Homing Adhesion Receptor; HCAM; HCELL; hematopoietic cell E-

Alternative Name: and L-selectin ligand; Heparan Sulfate Proteoglycan; Hermes Antigen; HAS; HUTCH I;

Hyaluronate Receptor; Indian blood group; Inlu Related p80 Glycoprotein; Ly 24; MDU2; MDU3;

MIC4; MUTCH I; Phagocytic Glycoprotein 1 (PGP-1)

Mouse IgG2a, kappa Isotype:

Immunogen Information: Recombinant fragment corresponding to the v6 domain of human CD44

Description

This antibody recognizes an epitope encoded by exon v6 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemopoiesis tissues. CD44 is a single gene with 20 exons, of which 10 are normally expressed to encode the basic CD44 (H-CAM) molecule. The additional 10 exons (v1 to v10) are only expressed by alternative splicing of the nuclear RNA. The expression of specific cell adhesion molecule CD44 splice variants has been reported to be associated with metastasis in certain human malignancies.

Product Info

Amount: $20 \mu g / 100 \mu g$

200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with Content:

0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is Storage condition:

stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Fig. 1: Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with CD44v6 Monoclonal Antibody (CD44v6/1246)



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Formalin-fixed, paraffin-embedded human Tongue Carcinoma stained with CD44v6 Monoclonal Antibody (CD44v6/1246)




Fig. 3: SDS-PAGE Analysis Purified CD44v6 Monoclonal Antibody (CD44v6/1246). Confirmation of Purity and Integrity of Antibody.