

36-3663: Anti-CD79b (B-Cell Marker) Monoclonal Antibody(Clone: rIGB/1842)

Clonality :	Monoclonal
Clone Name :	rlGB/1842
Application :	ELISA
Reactivity :	Human
Gene :	CD79B
Gene ID :	974
Uniprot ID :	P40259
Alternative Name :	AGM6; B-cell antigen receptor complex-associated protein beta chain; B-cell-specific glycoprotein B29; B29; B29/Ig-beta/CD79b; CD79b; Ig-beta; IGB; Immunoglobulin-associated B29 protein
Isotype :	Mouse IgG1, kappa
Immunogen Information	Recombinant human CD79b protein fragment (around aa29-159) (exact sequence is proprietary)

Description

CD79 (also designated Ig chains, designated CD79B or B29. The B cell antigen receptor complex (BCR) is formed by the association of CD79 with a membrane immunoglobulin, such as IgM or IgD. The membrane immunoglobulins IgM and IgD achieve surface expression and antigen presentation function in response to CD79 association. The cytoplasmic tails of both CD79A and CD79B contain an ITAM (immuno-receptor tyrosine-based activation) motif, which acts to initiate the BCR signaling reactions by binding to and activating tyrosine kinases.

Product Info	
Amount :	20 μg / 100 μg
Content :	200 μg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA);

kDa		R	NR	
250				
150 —			-	2ug loading NR=Non-
100 —				
75	-			
50	-		reduced	
37	-			R=reduced
²⁵	=	-	•	
15				
10	_			

Fig. 1: SDS-PAGE Analysis Purified CD79b Recombinant Mouse Monoclonal Antibody (rIGB/1842). Confirmation of Integrity and Purity of Antibody.