

### 36-3695: Anti-EBV Early Antigens (Epstein Barr Virus) Monoclonal Antibody(Clone: 1108-1)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	1108-1
<b>Application :</b>	IF
<b>Alternative Name :</b>	EBNA1; EBV nuclear antigen 1; Epstein Barr nuclear antigen 1; Human Herpesvirus 4 (HHV4)
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Affinity Purified early antigen polypeptides from induced Raji cells precipitated with African Burkitt's lymphoma sera

#### Description

Epstein-Barr virus (EBV), also designated human herpesvirus 4 (HHV-4), is a member of the herpesvirus family and is one of the most common human viruses. EBV infects B cells and, though often asymptomatic, it can cause infectious mononucleosis, a disease characterized by fatigue, fever, sore throat and muscle soreness. The EBV-induced early antigens (Ea) are among several antigen complexes that have been identified in EBV-infected cells. The Ea complex is composed of diffuse (Ea-D) and restricted (Ea-R) components. The activity of Ea-D is suppressed during latent infection. BMRF1, the gene that encodes for Ea-D, is closely associated with the gene encoding for EBV DNA polymerase, and Ea-D is essential for the activity of this polymerase. Ea-D forms a complex with EBV DNase and, together, they may play a role in viral replication.

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified EBV Mouse Monoclonal Antibody (1108-1). Confirmation of Purity and Integrity of Antibody.