

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3736: Anti-FITC (Fluorescein Isothiocyanate) Monoclonal Antibody(Clone: SPM395)

Clonality: Monoclonal Clone Name: SPM395

Application : Confocal Microscopy,FACS,IF,WB,IHC

Reactivity: All species

Alternative Name : Fluorescein isothiocyanate, FITC

Isotype: Mouse IgG1, kappa

Immunogen Information: Fluorescein isothiocyanate (FITC) conjµgated to KLH

Description

This MAb recognizes both the free and protein-conjµgated (either soluble or cell bound) form of fluorescein or carboxy-fluorescein. Sensitivity of the FITC-anti-FITC system is reportedly similar to that of the biotin-avidin system. In biotin or enzyme-labeled form, this MAb is also suitable for the conversation of fluorescein-labeled detection into immuno-enzymatic assays.

Product Info

Amount : $20 \mu g / 100 \mu g$

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 34 months. Non hazardous

is stable for 24 months. Non-hazardous.

Application Note

Confocal Microscopy; Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); In situ Nucleic Acid Hybridization; Western Blot; Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Pretreatment depends upon the fluoreceinated primary antibody.)