

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3803: Anti-Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) Monoclonal Antibody(Clone: KRTH/1576R)

Clonality: Monoclonal
Clone Name: KRTH/1576R

Application: IHC
Reactivity: Human
Gene: KRT76
Gene ID: 51350
Uniprot ID: Q01546

Alternative Name: KRT2B; KRT2P; HUMCYT2A; Keratin, type II Cytoskeletal 2 oral; K76; Keratin 2p (K2P);

Keratin-76; Cytokeratin-2P (CK-2P; Type-II Keratin Kb9

Isotype: Rabbit IgG

Immunogen Information: Recombinant full-length human KRT76 protein

Description

This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 52kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pl 6.0) subfamilies. The acidic keratins have molecular weights (MW) of 56.5, 55, 51, 50, 50', 48, 46, 45, and 40kDa. MAb AE3 recognizes the 65-67, 64, 59, 58, 56, and 52kDa keratins of basic subfamily. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis.

Product Info

Amount : $20 \mu g / 100 \mu g$

Content: 200 µg/ml of recombinant MAb Purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA

& 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Fig. 1: Formalin-fixed, paraffin-embedded human Skin stained with Cytokeratin, HMW Rabbit Recombinant Monoclonal Antibody (KRTH/1576R).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Formalin-fixed, paraffin-embedded human Skin stained with Cytokeratin, HMW Rabbit Recombinant Monoclonal Antibody (KRTH/1576R).




Fig. 3: SDS-PAGE Analysis Purified Cytokeratin, HMW Rabbit Recombinant Monoclonal Antibody (KRTH/1576R).