

### 36-3815: Anti-MECA-79 Monoclonal Antibody(Clone: MECA-79)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	MECA-79
<b>Application :</b>	ELISA,FACS,IF,WB,IHC
<b>Reactivity :</b>	Human, Mouse, Rat
<b>Alternative Name :</b>	MECA-79; Peripheral Node Addressin (PNAd)
<b>Isotype :</b>	Rat IgM, kappa
<b>Immunogen Information :</b>	Mouse lymph node stromal cells

#### Description

MECA-79 recognizes a carbohydrate epitope shared with a group of sulfation decorated sialomucins, including sulfated ligands for CD62L (CD34, GlyCAM-1, Sgp200, and a subset of MAdCAM-1). This set antigens has been referred to as peripheral node addressin (PNAd) with the molecular mass 50-250 kD. It has been identified that GlcNAc-6-SO<sub>4</sub> sulfation contributes to MECA-79 binding and the core 1 beta1,3-N-acetylglucosaminyltransferase is required for the generation of the MECA-79 epitope. MECA-79 is expressed on high endothelial venules (HEV) of lymphoid tissues, chronic inflamed tissues and rheumatoid synovia. The interaction of PNAd with CD62L receptor is involved in tethering and rolling of lymphocytes along HEV in lymphoid tissues. MECA-79 antibody reacts with Mouse, human and many other species PNAd and blocks L-selectin-dependent lymphocyte adhesion in vitro and in vivo.

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

ELISA (For coating, order antibody without BSA); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);