

### 36-3816: Anti-Human Herpes Virus 8 (HHV8) Monoclonal Antibody(Clone: LN53)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	LN53
<b>Application :</b>	ELISA,IF,IHC
<b>Alternative Name :</b>	HHV8; Human Herpes Virus 8; Human herpesvirus 8; Kaposi's sarcoma associated herpes virus; KSHV
<b>Isotype :</b>	Rat IgG2c, kappa
<b>Immunogen Information :</b>	Recombinant protein corresponding to the latent nuclear antigen 1 molecule of HHV8

#### Description

HHV 8 encodes a latent nuclear antigen (LNA), which is the product of the viral gene orf 73. LNA is capable of forming a complex with retinoblastoma susceptibility gene product, which may be related to its oncogenic activity. HHV8 is associated with three different diseases observed in AIDS patients; kaposi's sarcoma, primary effusion lymphoma (which is a rare type of non-Hodgkin lymphoma affecting the body cavities) and multicentric Castleman's disease. HHV 8 is the likely etiological agent of Kaposi's sarcoma.

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

ELISA (For coating use Ab at 1-5ug/ml, order Ab without BSA); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)




Fig. 1: SDS-PAGE Analysis Purified HHV8 Rat Monoclonal Antibody (LN53). Confirmation of Purity and Integrity of Antibody.