

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-190033: Recombinant Human ACE2 Protein with hFc tag

Application: **Functional Assay** 

Reactivity: Human 59272 Gene ID: **Uniprot ID:** Q9BYF1

Alternative Name: ACE-2; ACEH; ACE2

## **Description**

Source: HEK293 cells.

Endotoxin: < 0.1 EU/µg of the protein by LAL method.

Calculated MW: 109.5kDa

Recombinant Human ACE2 Protein is produced by HEK293 cells expression system. The target protein is expressed with

sequence (Gln18-Ser740) of human ACE2 (Accession #Q9BYF1) fused with a Fc tag at the C-terminus.

## **Product Info**

Amount: 100 µg

**Purification:** >95% by SDS-PAGE.

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. or Supplied as a 0.22 μm filtered Content:

solution in PBS, pH 7.4. Reconstitution: For lyophilized protein: Reconstitute to a concentration

of 0.1-0.5 mg/mL in sterile distilled water.

Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein

solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. This product is stable at ≤ Storage condition:

-70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.

Amino Acid: Recombinant Human ACE2 Protein is produced by HEK293 cells expression system. The target

protein is expressed with sequence (Gln18-Ser740) of human ACE2 (Accession #Q9BYF1) fused

with a Fc tag at the C-terminus.

## **Application Note**

Measured by its binding ability in a functional ELISA. Immobilized Recombinant 2019-nCoV Spike RBD-6His at 2μg/mL (100 μL/well) can bind Recombinant Human ACE2-Fc, the EC50 of ACE2-Fc is 4-6 ng/mL.




Fig 1: Recombinant Human ACE2 Protein with hFc tag was determined by SDS-PAGE with Coomassie Blue, showing a band at 140-150 kDa.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig 2 : Immobilized Recombinant 2019-nCoV Spike RBD-6His at  $2\hat{1}\frac{1}{4}$ g/mL (100  $\hat{1}\frac{1}{4}$ L/well) can bind Recombinant Human ACE2-Fc, the EC50 of ACE2-Fc is 4-6 ng/mL.