

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-4104: Phospho-MET(Tyr1234/1235) (Clone: 6F11) rabbit mAb APC conjugate

Clonality: Monoclonal

Clone Name: MetY12341235-6F11

Application: FACS

Reactivity: Human, Mouse

Conjugate: APC

Format: Conjugated

AUTS9; c-Met; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; HGFR; MET;

Alternative Name: met proto-oncogene (hepatocyte growth factor receptor); met proto-oncogene tyrosine kinase; oncogene MET; Proto-oncogene c-Met; RCCP2; Scatter factor receptor; SF receptor; Tyrosine-

protein kinase Met

Isotype: Rabbit IgG1k

Immunogen Information: A synthetic phospho-peptide corresponding to residues surrounding Tyr1234/Tyr1235 of human

phospho Met

Description

c-Met, also called tyrosine-protein kinase MET or hepatocyte growth factor receptor (HGFR), has tyrosine kinase activity (1). MET is a single pass tyrosine kinase receptor essential for embryonic development, organogenesis and wound healing. Normally, MET is expressed only in stems cells and progenitor cells but excessive expression of MET/HGFR and its autocrine activation by co-expression of hepatocyte growth factor (HGF) ligand are implicated in oncogenesis (2,3). Aberrantly activated MET leads to tumor growth, angiogenesis, and cancer metastasis and is correlated with poor prognosis. Abnormal activation of MET is observed in various human malignancies, such as kidney, liver, stomach, breast, and brain. MET activation by HGF induces MET kinase catalytic activity and leads to phosphorylation at Tyr 1234 and Tyr 1235.

Product Info

Amount: 10 Tests / 100 Tests

Content: 1X PBS, 0.09% NaN3, 0.2% BSA

Storage condition : Store at 2-8°C. Avoid repeated freeze and thaw cycles.

Application Note

For flow cytometric staining, the suggested use of this reagent is 5 $\text{Å}\mu\text{L}$ per million cells or 5 $\text{Å}\mu\text{L}$ per 100 $\text{Å}\mu\text{L}$ of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.




Fig-1: Flow cytometric analysis of Ramos cells unstained untreated cells (blue) or stained untreated (red) or treated with FBS and pervanadate (green) using phospho-MET(Y1234/1235) antibody METY12341235-6F11 APC conjugate.