# **w** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

## 12-4113: Phospho-Ship2 (Tyr1135) (Clone: 1D2) rabbit mAb APC conjugate

Clonality :	Monoclonal
Clone Name :	Ship2Y1135-1D2
Application :	FACS
Reactivity :	Human, Mouse
Conjugate :	APC
Format :	Conjugated
Alternative Name :	Phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase 2, Inositol polyphosphate phosphatase-like protein 1, INPPL-1, Protein 51C, SH2 domain-containing inositol 5'-phosphatase 2
Isotype :	Rabbit lgG1k
Immunogen Information	A synthetic phospho-peptide corresponding to residues surrounding Tyr1135 of human phospho Ship2

#### Description

Ship2 (SH2-containing inositol phosphatase 2, phospho Ship2) is a homolog of Ship1. Ship2 is highly expressed in the heart, in skeletal muscle, and in the placenta (1). SHIP2 negatively regulates insulin signaling (2), and Ship2 polymorphisms have been linked to hyperglycemia (3). Phospho Ship2 has been identified as a potential therapeutic target for the treatment of obesity and type 2 diabetes (4,5). The Ship2 residue Tyr1135 is phosphorylated in human cancer cells (6-8).

#### **Product Info**

Amount :	10 Tests / 100 Tests
Content :	1X PBS, 0.09% NaN3, 0.2% BSA
Storage condition :	Store at 2-8°C. Avoid repeated freeze and thaw cycles.

### **Application Note**

For flow cytometric staining, the suggested use of this reagent is 5  $\tilde{A}$   $\hat{A}\mu$  per million cells or 5  $\tilde{A}$   $\hat{A}\mu$  per 100  $\tilde{A}\mu$  of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.




Fig-1: Flow cytometric analysis of U937 cells secondary antibody only negative control (blue) or untreated (red) or treated with IFNa IL-4 and pervanadate (green) using Phospho-Ship2 (Tyr1135) APC conjugated antibody Ship2Y1135-1D2.