

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-4114: Phospho-Ship2 (Tyr1135) (Clone: 1D2) rabbit mAb SureLight488 conjugate

Clonality: Monoclonal
Clone Name: Ship2Y1135-1D2

Application: FACS

Reactivity: Human, Mouse
Conjugate: SureLight 488
Format: Conjugated

Phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase 2, Inositol polyphosphate

Alternative Name: phosphatase-like protein 1, INPPL-1, Protein 51C, SH2 domain-containing inositol 5'-

phosphatase 2

Isotype: Rabbit IgG1k

Immunogen Information: A synthetic phospho-peptide corresponding to residues surrounding Tyr1135 of human

phospho Ship2

Description

Ship2 (SH2-containing inositol phosphatase 2, phospho Ship2) is a homolog of Ship1. Ship2 is highly expressed in the heart, in skeletal muscle, and in the placenta (1). SHIP2 negatively regulates insulin signaling (2), and Ship2 polymorphisms have been linked to hyperglycemia (3). Phospho Ship2 has been identified as a potential therapeutic target for the treatment of obesity and type 2 diabetes (4,5). The Ship2 residue Tyr1135 is phosphorylated in human cancer cells (6-8).

Product Info

Amount: 10 Tests / 100 Tests

Content: 1X PBS, 0.09% NaN3, 0.2% BSA

Storage condition : Store at 2-8°C. Avoid repeated freeze and thaw cycles.

Application Note

For flow cytometric staining, the suggested use of this reagent is $5 \text{ A} \Box \hat{A} \mu L$ per million cells or $5 \text{ A} \Box \hat{A} \mu L$ per 100 $\hat{A} \Box \hat{A} \mu L$ of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.




Fig-1: Flow cytometric analysis of U937 cells unstained untreated U937 cells as negative control (blue) or stained and untreated (red) or treated with IFNa IL-4 and pervanadate (green) using phospho-Ship2 (Tyr1135) S antibody Ship2Y1135-1D2 ureLight 488 conjugated.