

12-4116: Phospho-PTEN (Ser380) (Clone: NA9) rabbit mAb

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | PTENS380-NA9 |
| Application : | FACS |
| Reactivity : | Human, Mouse |
| Conjugate : | Unconjugated |
| Format : | Purified |
| Alternative Name : | Phosphatidylinositol 3,4,5-trisphosphate 3-phosphatase and dual-specificity protein phosphatase, Mutated in multiple advanced cancers 1, MMAC1, Phosphatase and tensin homolog, TEP1 |
| Isotype : | Rabbit IgG1k |
| Immunogen Information : | A synthetic phospho-peptide corresponding to residues surrounding Ser380 of human phospho PTEN |

Description

PTEN has been identified as a tumor suppressor gene and has been found to be mutated in a significant number of human cancers, including prostate, brain, and breast cancer. PTEN shares sequence homology with the protein-tyrosine phosphatase (PTPase) family of proteins and negatively regulates the PI3K/Akt pathway. PTEN de-phosphorylates target proteins, and recombinant PTEN has been shown to have phosphoinositide 3-phosphatase and inositol phosphate 3-phosphatase activity. Studies of primary tumor cells show a loss of PTEN expression after metastasis to the brain, via astrocyte-derived microRNAs. A cluster of phosphorylation sites (S380, T382, T383, and S385) in the C-terminal tail of PTEN drive a conformational change that reduces PTEN activity by inhibiting membrane interactions.

Product Info

| | |
|----------------------------|---|
| Amount : | 20 µl / 200 µl |
| Content : | 1X PBS, 0.02% NaN ₃ , 50% Glycerol, 0.1% BSA |
| Storage condition : | Store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

1Âµg/mL - 0.001Âµg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.(0.5mg/ml)




Fig-1: Flow cytometric analysis of A431 cells, untreated and unstained as negative control (blue) or untreated and stained (green) or treated with lambda phosphatase and stained (red) using Phospho-PTEN (S380) antibody, PTENS380-NA9 at 0.1 µg/mL.




Fig 2 : Peptide blocking flow cytometric analysis of A431 cells secondary antibody only negative control (light blue) or untreated (red) or treated with EGF (green) or untreated and blocked with phospho-peptide (black) or EGF and blocked with phospho peptide (gold) or untreated and blocked with non-phospho peptide (dark blue) or EGF and blocked with non-phospho peptide (purple) using Phospho-PTEN (S380) antibody PTENS380-NA9 0.05 µg/mL.




Fig-3: PTENS380-NA9 recognizes basal phosphorylation levels in mouse cells. Flow cytometric analysis of L929 cells secondary antibody only (blue) or 0.1 µg/mL of isotype control (orange) or of Phospho-PTEN (S380) antibody PTENS380-NA9 (green).