

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-4148: Phospho-Zap70 (Tyr493)/Syk (Tyr526) (Clone: H11) rabbit mAb APC conjugate

Clonality: Monoclonal
Clone Name: Zap70Y493-H11

Application :FACSReactivity :HumanConjugate :APC

Alternative Name:

Tyrosine-protein kinase ZAP-70 , 70 kDa zeta-chain associated protein, Syk-related tyrosine

kinase, SRK, Tyrosine-protein kinase SYK, Spleen tyrosine kinase, p72-Syk

**Isotype:** Rabbit IgG1k

**Immunogen Information**: A synthetic phospho-peptide corresponding to residues surrounding Tyr493 of human

phospho Zap70

## **Description**

ZAP70 (Tyrosine-protein kinase ZAP-70, phospho Zap70) is a protein tyrosine kinase (PTK) that associates with the z subunit of the T cell antigen receptor (TCR) and undergoes tyrosine phosphorylation following TCR stimulation. Following TCR engagement, Zap-70 is rapidly phosphorylated on several tyrosine residues through autophosphorylation and transphosphorylation by the Src family tyrosine kinase Lck. ZAP70 contains two SH2-like domains with the PTK domain located at the C-terminus. It appears that both phospho Zap70 and Syk are recruited to the phosphorylated CD3 and z subunits after TCR stimulation. Phosphorylation of Tyr319 is required for the assembly of a phospho Zap70-containing signaling complex that leads to the activation of the PLC-gamma1-dependent and Ras-dependent signaling cascades in antigen-stimulated T cells. The orthologous Tyr352 residue in Syk is also involved in the association with PLC-gamma1.

## **Product Info**

Amount: 10 Tests / 100 Tests

**Content:** 1X PBS, 0.09% NaN3, 0.2% BSA

**Storage condition :** Store at 2-8°C. Avoid repeated freeze and thaw cycles.

## **Application Note**

For flow cytometric staining, the suggested use of this reagent is 5  $\mu$ L per million cells or 5  $\mu$ L per 100  $\mu$ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.




Fig-1: Flow cytometric analysis of Jurkat cells unstained cells negative control (blue) or stained and untreated (red) or treated with pervanadate (green) using Phospho-Zap70 (Tyr493)/Syk (Tyr526) antibody Zap70Y493-H11-APC conjugate.