

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-4195: Phospho-MSK1 (Thr581) (Clone: A5) rabbit mAb PE conjugate

Clonality: Monoclonal
Clone Name: MSK1T581-A5

Application: FACS

Reactivity: Human, Mouse, Rat

Conjugate: PE

Format: Conjugated

Ribosomal protein S6 kinase alpha-5, S6K-alpha-5, 90 kDa ribosomal protein S6 kinase 5,

Alternative Name: Nuclear mitogen- and stress-activated protein kinase 1, RSK-like protein kinase, RSKL,

RPS6KA5

Isotype: Rabbit IgG1k

Immunogen Information: A synthetic phospho-peptide corresponding to residues surrounding Thr581 of human

phospho MSK1

Description

MSK1 (mitogen and stress activated protein kinase 1, phospho MSK1) is activated by Erk in response to growth factors and by and p38 in response to cellular stress (1). MSK1 is similar to RSK1 in that it has two kinase domains and a connecting regulatory linker region (2). S364/S381 phosphorylation activates RSK1 (3), which is analogous to residues S360 and S376 of MSK1, which may be important for phospho MSK1 activity.

Product Info

Amount: 10 Tests / 100 Tests

Content: 1X PBS, 0.09% NaN3, 0.2% BSA

Storage condition : Store at 2-8°C. Avoid repeated freeze and thaw cycles.

Application Note

For flow cytometric staining, the suggested use of this reagent is $5 \text{ A} \triangle \mu$ per million cells or $5 \text{ A} \triangle \mu$ per 100 A $\triangle \mu$ of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.




Fig-1: Flow cytometric analysis of COS7 cells unstained and untreated (blue) or untreated (red) or treated with TPA + CalA (green) and stained using phospho-MSK1(T581) PE conjugated antibody MSK1T581-A5.