

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

12-4244: Phospho-Lck (Tyr505) (Clone: A3) rabbit mAb FITC conjugate

Clonality: Monoclonal LckY505-A3 Clone Name: Application: **FACS** Human Reactivity: Conjugate: **FITC**

Format: Conjugated

Tyrosine-protein kinase Lck, Leukocyte C-terminal Src kinase, LSK, Protein YT16, T cell-specific **Alternative Name:**

protein-tyrosine kinase

Rabbit IgG1k Isotype:

A synthetic phospho-peptide corresponding to residues surrounding Tyr505 of human phospho Immunogen Information:

Description

Lck is a member of the Src family of non-receptor tyrosine kinases and plays a major role in T cell activation. Lck activates many downstream signaling pathways including Akt/mTOR, SAPK/JNK, PLCg1, and RAS/MAPK. Phosphorylation of Lck at Tyr394 in the catalytic domain at the ATP-binding site stabilizes the open and active form, while phosphorylation at Tyr505 in the Cterminal domain promotes the closed, inactive conformation. Multiple small-molecule drugs used to treat leukemia have been shown to target inhibition of Lck, including imatinib and dasatinib. Lck is thus a promising target for suppressing T-cell responses for the treatment of inflammatory diseases or after organ transplantation.

Product Info

Amount: 10 Tests / 100 Tests

Content: 1X PBS, 0.09% NaN3, 0.2% BSA

Store at 2-8°C. Avoid repeated freeze and thaw cycles. Storage condition:

Application Note

For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells or 5 µL per 100 µL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.




Fig-1: Flow cytometric analysis of Daudi Human Burkitt's lymphoma cells untreated and unstained as negative control (blue) or untreated (red) or treated with IFNa +IL4 +PV and stained (green) using Phospho-LCK (Y505) antibody LCKY505-A3 FITC conjugate.