

12-4272: Phospho-Btk (Tyr551) (Clone: G12) rabbit mAb

Clonality :	Monoclonal
Clone Name :	BtkY551-G12
Application :	FACS,ELISA
Reactivity :	Human
Conjugate :	Unconjugated
Format :	Purified
Alternative Name :	Bruton tyrosine kinase, Tyrosine-protein kinase BTK, Agammaglobulinemia tyrosine kinase, ATK, AGMX1, B-cell progenitor kinase, BPK
Isotype :	Rabbit IgG1k
Immunogen Information :	A synthetic phospho-peptide corresponding to residues surrounding Tyr551 of human phospho Btk

Description

Btk is a major node in the B-cell receptor signaling pathway, where it regulates B cell maturation, activation, survival, differentiation, and proliferation. Btk is activated by Src family kinases, including Lyn, which phosphorylates Btk at Tyr551. Upon phosphorylation at this site, Btk is recruited to the plasma membrane where autophosphorylation at Tyr223 occurs. The Btk signaling pathway is a major target of small molecule inhibitors for B-cell lymphoma, autoimmune diseases, and non-Hodgkin's lymphomas. These inhibitors either form a covalent bond at Cys481 in the ATP-binding site or serve as reversible inhibitors that bind the SH3 pocket and stabilize inactive Btk.

Product Info

Amount :	20 µl / 200 µl
Content :	1X PBS, 0.02% NaN ₃ , 50% Glycerol, 0.1% BSA
Storage condition :	Store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

1 µg/mL - 0.001 µg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.(0.5mg/ml, more than 200 western blots)




Fig-1: Flow cytometric analysis of Ramos cells secondary antibody only negative control (blue) or untreated (red) or treated with pervanadate (green) using Phospho-Btk (Tyr551) antibody BtkY551-G12 at 0.1 µg/mL.




Fig 2 : Flow cytometric analysis of Ramos cells secondary antibody only negative control (blue) or treated with imatinib (red) or with pervanadate (green) using Phospho-Btk (Y551) antibody BtkY551-G12 or Company B antibody at 0.1 µg/mL (manufacturer's recommended concentration).




Fig-3: Peptide ELISA using BtkY551-G12 titrated starting from 0.3 µg/mL shows binding to only Btk pY551 phospho peptide and no cross-reactivity to Btk Y551 non-phospho peptide or to control peptides.




Fig-4: Direct ELISA using Ramos cellular lysate coated directly to the plate surface after lysis following no treatment or treatment with 1mM pervanadate for 30 min. ELISA wells were tested in duplicate using 0.25 mg/mL total protein coated lysate and 1 µg/mL BtkY551-G12 IgG.