

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-4317: Phospho-NFkB p65 (Ser529) (Clone: H3) rabbit mAb PE conjugate

Clonality: Monoclonal

Clone Name: NFkBp65S529-H3

Application: FACS

Reactivity: Human, Mouse

Conjugate: PE

Format: Conjugated

Alternative Name: Transcription factor p65, Nuclear factor NF-kappa-B p65 subunit, Nuclear factor of kappa light

polypeptide gene enhancer in B-cells 3, RELA, NFKB3

Isotype: Rabbit IgG1k

Immunogen Information : A synthetic phospho-peptide corresponding to residues surrounding Ser529 of human phospho-

NFKB p65

## **Description**

The nuclear factor kB (NFkB)/Rel family of transcription factors play a pivotal role in inflammatory and immune responses (1,2). NF-kappa-B is present in almost all cell types and is involved in many biological processes including immunity, inflammation, cell growth and differentiation, apoptosis, and tumorigenesis. NFkB is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFkB1/p105, NFkB1/p50, REL and NFkB2/p52. The dimers bind at kB sites in the target gene DNA. Individual dimers have distinct preferences for different kB sites and can act as either transcriptional activators or repressors. NFkB Ser536 phosphorylation stimulates Lys310 acetylation and interaction of phospho NFkB with CBP. Acetylated/phospho NFkB induces enhanced transcriptional activity.

## **Product Info**

Amount: 10 Tests / 100 Tests

Content: 1X PBS, 0.09% NaN3, 0.2% BSA

**Storage condition :** Store at 2-8°C. Avoid repeated freeze and thaw cycles.

## **Application Note**

For flow cytometric staining, the suggested use of this reagent is 5  $\text{\^A}\mu\text{L}$  per million cells or 5  $\text{\^A}\mu\text{L}$  per 100  $\text{\^A}\mu\text{L}$  of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.




Fig-1: Flow cytometric analysis of HeLa cells untreated and unstained as negative control (blue) or untreated (red) or treated with TNFa + calyculin A (green) and stained using Phospho-NFkB p65 (Ser529) antibodyPE conjugate, NFkBp65S529-H3.