

12-4334: Phospho-Ask1 (Thr838) (Clone: 8D12) rabbit mAb

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | Ask1T838-8D12 |
| Application : | FACS |
| Reactivity : | Human, Mouse |
| Conjugate : | Unconjugated |
| Format : | Purified |
| Alternative Name : | Apoptosis signal-regulating kinase 1, Mitogen-activated protein kinase kinase kinase 5, MAPK/ERK kinase kinase 5, MAP3K5, MAPKKK5, MEKK5 |
| Isotype : | Rabbit IgG1k |
| Immunogen Information : | A synthetic phospho-peptide corresponding to residues surrounding Thr838 of human phospho Ask1 |

Description

Stress signaling in cells are partly mediated through c-Jun N-terminal kinase (JNK) and p38 mitogen-activated protein kinase (MAPK). A kinase upstream of these stress-mediated pathways is apoptosis signal-regulating kinase 1 (ASK1). ASK1 is a serine/threonine kinase that belongs to the family of mitogen-activated protein kinase kinase kinase (MAP3K). It is endogenously expressed in various cells. Various stress signaling activates ASK1 including reactive oxygen species (ROS), tumor necrosis factor alpha (TNFα), lipopolysaccharide (LPS) and endoplasmic reticulum (ER) stress. These events ultimately lead to the activation of JNK and p38 MAPK. As a result, ASK1-MAPK signaling is associated with various cellular stress responses including apoptosis, differentiation, and inflammation. Several studies show that aberrant ASK1 signaling leads to severe human diseases such as neurodegenerative diseases, induction of inflammatory diseases and cancer. The phosphorylation of the threonine residues 838 (Thr838) in human and Thr845 in mice is important for ASK1 activation. At basal inactive state, ASK1 is a homooligomer, which binds to another ASK1 via its C-terminal coiled-coil domain. The N-terminal coiled-coil of ASK1 binds to thioredoxin (Trx), which suppresses ASK1 kinase activity. Under oxidative stress, oxidized Trx is separated from ASK1 and unbound ASK1 is activated by phosphorylation. Calcium influx and oxidative stress can elicit phosphorylation of the ASK1. After H₂O₂ injury, tumor necrosis factor 2, (TRAF2) and TRAF6 act as positive regulator of ASK1. 14-3-3 proteins, act as negative regulator of ASK1 by binding to the C-terminal of ASK1 after Ser966 phosphorylation. However oxidative stress promotes dephosphorylation of Ser966 and lead to detachment of 14-3-3. TNF and Fas death receptor activate ASK1. ASK1 is an early signal stress responder to oxidative stress in cerebral ischemia. ASK1 has also been implicated in thrombosis, brain edema, inflammatory response, and reactive gliosis.

Product Info

| | |
|----------------------------|---|
| Amount : | 20 µl / 200 µl |
| Content : | 1X PBS, 0.02% NaN ₃ , 50% Glycerol, 0.1% BSA |
| Storage condition : | Store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

1 Âµg/mL - 0.001 Âµg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.(0.5mg/ml, more than 200 western blots)




Fig-1: Flow cytometric analysis of NIH3T3 cells treated with staurosporine and unstained as negative control (blue) or treated with staurosporine (red) or with PDGF (green) and stained using Phospho-Ask1 (Thr838) antibody Ask1T838-8D12 at 5ng/mL.




Fig 2 : Peptide blocking flow cytometric analysis of 3T3 cells secondary antibody only negative control (light blue) or staurosporine-treated (red) or PDGF-treated (green) or staurosporine and blocked with phospho-peptide (black) or PDGF and blocked with phospho peptide (gold) or staurosporine and blocked with non-phospho peptide (dark blue) or PDGF and blocked with non-phospho peptide (purple) using Ask1 (Thr838) antibody Ask1T838-8D12 at 10ng/mL.