

## 32-9190: Recombinant Mouse FCRN Heterodimer

**Alternative Name :** IgG receptor FcRn; Neonatal Fc receptor; FCRN; FCGRT&B2M

### Description

Source : Human 293 Cells;

FCGRT & B2M heterodimer protein (FcRn complex) consist of two subunits: p51, also known as FCGRT, and p14, also known as beta-2-microglobulin, and forms an MHC class I-like heterodimer. It is widely expressed in endothelial and epithelial cells and plays an important role in IgG homeostasis. Fc fragment of IgG, receptor, transporter, alpha (FCGRT) binds to the Fc region of monomeric immunoglobulins gamma and mediates the uptake of IgG from milk. FCGRT thus play a role in the transfer of immunoglobulin G from mother to fetus. B2M is a component of class I major histocompatibility complex (MHC) and involved in the presentation of peptide antigens to the immune system.

### Product Info

**Amount :** 500 µg / 50 µg

**Content :** Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4

**Amino Acid :** Recombinant Mouse IgG Fc fragment receptor transporter is produced by our Mammalian expression system and the target gene encoding Ser22-Ser297&Ile21-Met119 is expressed with a 6His tag at the C-terminus.