

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-9359: Recombinant Mouse Tacstd2/Trop2(C-6His)

Alternative Name: Tumor-associated calcium signal transducer 2;Tacstd2;Trop2;Cell surface glycoprotein Trop-2

## **Description**

Source: Human Cells;

Tumor-associated calcium signal transducer 2(Tacstd2), also known as Cell surface glycoprotein Trop-2, belongs to the EPCAM family. Tacstd2 expressed in kidney, lung, ovary and testis and has high levels of expression in immortalized keratinocytes. Tacstd2 may functions as a growth factor receptor. It has negative regulation of branching involved in ureteric bud morphogenesis, cell motility, epithelial cell migration, ruffle assembly, stress fiber assembly and substrate adhesion-dependent cell spreading. Also, it has positive regulation of stem cell proliferation. Tacstd2 is capable of transducing an intracellular calcium signal and may play a role in tumor growth. It also has adhesive functions.

## **Product Info**

**Amount:** 500 μg / 50 μg

 $\textbf{Content}: \ \, \text{Lyophilized from a 0.2} \ \, \mu\text{m filtered solution of PBS,pH7.4}.$ 

Amino Acid: Recombinant Mouse Tumor-associated calcium signal transducer 2 is produced by our Mammalian

expression system and the target gene encoding Gln25-Gly270 is expressed with a 6His tag at the

C-terminus.