

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-9477: Recombinant Human Interleukin-12 Subunit beta/IL-12 p40/IL-12B

Alternative Name: Interleukin-12 subunit beta, IL-12B, Cytotoxic lymphocyte maturation factor 40 kDa subunit, CLMF p40, IL-12 subunit p40, NK cell stimulatory factor chain 2, NKSF2, IL12B, NKSF2

Description

Source: Human Cells;

Interleukin-12 subunit beta (IL-12B) belongs to the type I cytokine receptor family. It contains 1 fibronectin type-III domain and 1 Ig-like C2-type domain. IL-12B is a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. IL-12 is a disulfide-linked heterodimer composed of the 40 kD cytokine receptor encoded by IL12B and a 35 kD subunit encoded by IL12A. IL12 is expressed by activated macrophages that serve as an essential inducer of Th1 cells development. It has been found to be important for sustaining a sufficient number of memory/effector Th1 cells to mediate long-term protection to an intracellular pathogen.

Product Info

Amount : 500 μg / 50 μg

Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Amino Acid: Recombinant Human Interleukin-12 subunit beta is produced by our Mammalian expression

system and the target gene encoding Ile23-Ser328 is expressed.