

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-9639: Recombinant Human TIGIT/VSIG9/VSTM3 (C-mFC)

Alternative Name : T-cell immunoreceptor with Ig and ITIM domains;;VSIG9; VSTM3;TIGIT;V-set and transmembrane domain-containing protein 3;V-set and immunoglobulin domain-containing protein 9

Description

Source: Human Cells;

T cell immunoreceptor with Ig and ITIM domains (TIGIT) is a member of the CD28 family within the Ig superfamily of proteins. TIGIT is expressed on NK cells and subsets of activated, memory and regulatory T cells, and particularly on follicular helper T cells within secondary lymphoid organs. It binds to CD155 and Nectin-2 that appear on dendritic cells (DC) and endothelium. Ligation of TIGIT on T cells down-regulates TCR-mediated activation and subsequent proliferation, while NK cell TIGIT ligation blocks NK cell cytotoxicity. Through CD155 and Nectin-2, which also interact with DNAM-1/CD226 and CD96/Tactile, TIGIT is part of an interacting network of Ig superfamily members that may augment or oppose each other. In particular, TIGIT binding to CD155 can antagonize the effects of DNAM1.

Product Info

Amount : $500 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Amino Acid: Recombinant Human T Cell Immunoreceptor With Ig And ITIM Domains is produced by our

Mammalian expression system and the target gene encoding Met22-Pro141 is expressed with a

mFc tag at the C-terminus.