

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-9686: Recombinant Human OX-2/MOX1/CD200 (C-Fc)

Alternative Name: OX-2 Membrane Glycoprotein, CD200, MOX1, MOX2

## **Description**

Source: Human Cells;

CD200 is a transmembrane immunoregulatory protein that belongs to the immunoglobulin superfamily. It contains one Ig like V type domain and one Ig like C2 type domain in its extracelluar domain. CD200 is widely but not ubiquitously expressed. Its receptor (CD200R) is restricted primarily to mast cells, basophils, macrophages, and dendritic cells, which suggests myeloid cell regulation as the major function of CD200. CD200 and CD200R associate via their respective N-terminal Ig-like domains. In myeloid cells, CD200R initiates inhibitory signals following receptor-ligand contact. In T cells, CD200 functions as a costimulatory molecule independent of the CD28 pathway. In addition, CD200 also plays an important role in prevention of graft rejection, autoimmune diseases and spontaneous abortion.

## **Product Info**

**Amount**: 500 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Amino Acid: Recombinant Human CD200 is produced by our Mammalian expression system and the target

gene encoding Gln31-Gly232 is expressed with a Fc tag at the C-terminus.