

## 32-6425: IL-4 Porcine

**Application :** Functional Assay  
**Alternative Name :** BCGF, BCDF, B cell stimulating factor, BSF-1, Lymphocyte stimulatory factor 1, IL-4, MGC79402, Binetrakin, Pitrakinra.

### Description

Source: Escherichia Coli.

Sterile Filtered White lyophilized (freeze-dried) powder.

IL4 is a pleiotropic cytokine produced by activated T cells. IL4 is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which contributes to various overlapping roles of this cytokine and IL13. STAT6, a signal transducer and activator of transcription plays a main role in mediating the immune regulatory signal of this cytokine. IL4, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. IL4, IL13 and IL5 are regulated co-ordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome.

Interleukin-4 Porcine Recombinant produced in E. coli is a non-glycosylated monomer chain containing 110 amino acids and having a molecular mass of 12.7kDa. IL-4 is purified by proprietary chromatographic techniques.

### Product Info

**Amount :** 50 µg / 100 µg  
**Purification :** Greater than 95.0% as determined by SDS-PAGE.  
**Content :** The protein was lyophilized from a sterile (0.2µm) filtered solution containing 10 mM sodium phosphate, pH 7.5.  
It is recommended to reconstitute the lyophilized IL-4 in sterile 18M Omega -cm H<sub>2</sub>O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.  
**Storage condition :** Lyophilized IL-4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL-4 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.  
**Amino Acid :** MHKCDITLQE IIKTLNILTA RKNSCMELPV TDVFAAPENT TEKETF CRAS TVLRHIYRHH  
TCMKLLSGL DRNLSSMANM TCSVHEAKKS TLKDFLERLK TIMKEKYSKC.

### Application Note

The ED<sub>50</sub>, as determined by TF-1 cell proliferation is 0.504ng/ml corresponding to a specific activity which is 2.0 x 10<sup>6</sup> units/mg.