

## 32-6473: IL 29 Human, Sf9

**Alternative Name :** Interleukin-29, IL-29, IFN-Lambda 1, Interferon-Lambda 1, Cytokine ZCYTO21, IL29, IFNL1, ZCYTO21.

### Description

Source: Sf9, Baculovirus cells.

Sterile filtered colorless solution.

IL-29 is distantly related to type I interferons and the IL-10 family. Expression of IL-29 is induced by viral infection which interacts with a heterodimeric class II cytokine receptor that consists of interleukin 10 receptor, beta (IL10RB) and interleukin 28 receptor, alpha. IL-29 exhibits common features with type I IFNs such as antiviral activity, antiproliferative activity and in vivo antitumour activity. IL-29 acts similarly to IFNs, but is less effective generally and has activity in a more limited range of cell lines. IFN-ambda 1, IFN-lambda 2 and IFN-lambda3 are closely positioned genes on human chromosome 19. IL-29 induces ELR(-) CXC chemokine mRNA in human peripheral blood mononuclear cells, in an IFN-gamma-independent manner. IL-29 is able to generate tolerogenic DCs, an activity that could thwart IFN-beta functions. IL-29 produced in response to viral infection, activates both monocytes and macrophages producing a restricted panel of cytokines and therefore is an important factor in activating innate immune responses at the site of viral infection. IFN-Lambda 1 antiviral and antiproliferative activity requires Interferon-Lambda 2 receptor tyrosine residues.

Interleukin-29 Human Recombinant produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 187 amino acids (20-200a.a.) and having a molecular mass of 20.8kDa (Molecular size on SDS-PAGE will appear at approximately 18-28kDa). IL29 is fused with a 6 amino acids His tag at C-Terminus and purified by proprietary chromatographic techniques.

### Product Info

**Amount :** 2 µg / 10 µg

**Purification :** Greater than 90.0% as determined by SDS-PAGE.

**Content :** IL29 protein solution (0.5mg/ml) contains Phosphate Buffered Saline (pH7.4) and 10% glycerol.

**Storage condition :** Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

**Amino Acid :** GPVPTSKPTT TGKGCHIGRF KSLSPQELAS FKKARDALEE SLKLKNWSCS SPVFPGNWDL RLLQVRERPV  
ALEAELALT LKLEAAAGPA LEDVLDQLH TLHHLSQLQ ACIQPQPTAG PRPRGRLHHW LHRLQEAPKK  
ESAGCLEASV TFNLFRLLTR DLKYVADGNL CLRTSTHPES THHHHHH.