

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-6522: PLGF1 Human, 132 a.a.

Application : Functional Assay **Alternative Name :** PIGF, PGF, PLGF-1.

Description

Source: Escherichia Coli.

Sterile Filtered White lyophilized (freeze-dried) powder.

PLGF is a growth factor active in angiogenesis, and endothelial cell growth, stimulating their proliferation and migration. It binds to receptor vegfr-1/flt1.

Placental Growth Factor-1 Human Recombinant produced in E.Coli is a non-glycosylated, disulfide-linked homodimer consisting of 2x132 amino acid polypeptide chains, having a total molecular mass of approximately 29.7kDa. PLGF1 is purified by proprietary chromatographic techniques.

Product Info

Amount: $2 \mu g / 10 \mu g$

Purification: Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Lyophilized from a $0.2\mu m$ filtered concentrated solution in 10mM Sodium Phosphate pH 7.5. It is recommended to reconstitute the lyophilized Placental Growth Factor-1 in sterile 18M

Content:

Omega -cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous

solutions.

Lyophilized PLGF1 although stable at room temperature for 3 weeks, should be stored

Storage condition: desiccated below -18°C. Upon reconstitution Placental Growth Factor-1 should be stored at 4°C

between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Amino Acid: MLPAVPPQQW ALSAGNGSSE VEVVPFQEVW GRSYCRALER LVDVVSEYPS EVEHMFSPSC

VSLLRCTGCC GDENLHCVPV ETANVTMQLL KIRSGDRPSY VELTFSQHVR CECRPLREKM KPERCGDAVP

RR.

Application Note

Determined by its ability to chemoattract human monocytes of using a concentration range of 1.0-10.0 ng/ml.