

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-6587: TPO Human, His

MPL Ligand, THCYT1, MPLLG, TPO, ML, Thrombopoietin Nirs Variant 1, MKCSF, Thrombopoietin,

Alternative Myeloproliferative Leukemia Virus Oncogene Ligand, Megakaryocyte Growth and Development Factor,
Name: Megakaryocyte Colony-Stimulating Factor, C-Mpl Ligand, MGDF, Megakaryocyte Stimulating Factor, Prepro-

Thrombopoietin.

Description

Source: Sf9, Insect cell. Sterile Filtered clear solution.

TPO produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 340 amino acids (22-353aa.) and having a molecular mass of 36.5kDa. (Molecular size on SDS-PAGE will appear at approximately 40-57KDa). TPO is expressed with an 8 amino acid His tag at C-Terminus and purified by proprietary chromatographic techniques..

Product Info

Amount: $2 \mu g / 10 \mu g$

Purification: Greater than 85% as determined by SDS-PAGE.

Content: TPO protein solution (0.25mg/ml) contains phosphate buffered saline (pH7.4) and 10% glycerol.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of

Storage condition : time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid

multiple freeze-thaw cycles.

Amino Acid: SPAPPACDLR VLSKLLRDSH VLHSRLSQCP EVHPLPTPVL LPAVDFSLGE WKTQMEETKA

QDILGAVTLL LEGVMAARGQ LGPTCLSSLL GQLSGQVRLL LGALQSLLGT QLPPQGRTTA

HKDPNAIFLS FQHLLRGKVR FLMLVGGSTL CVRRAPPTTA VPSRTSLVLT

LNELPNRTSGLLETNFTASA RTTGSGLLKW QQGFRAKIPG LLNQTSRSLD QIPGYLNRIH ELLNGTRGLF PGPSRRTLGA PDISSGTSDT GSLPPNLQPG YSPSPTHPPT GQYTLFPLPP

TLPTPVVQLH PLLPDPSAPT PTPTSPLLNT SYTHSQNLSQ EGLEHHHHHH.