

32-6605: Fractalkine Human, Sf9

Alternative Name : Fractalkine, CX3CL1, Neurotactin, CX3C membrane-anchored chemokine, Small inducible cytokine D1, NTN, NTT, CXC3, CXC3C, SCYD1, ABCD-3, C3Xkine.

Description

Source: Sf9, Baculovirus cells.

Sterile Filtered colorless solution.

Fractalkine soluble form is chemotactic for t-cells and monocytes, but not for neutrophils. Fractalkine membrane-bound form promotes adhesion of those leukocytes to endothelial cells. Fractalkine regulates leukocyte adhesion and migration processes at the endothelium and binds to CX3CR1. Fractalkine gene is located on human chromosome 16 along with some CC chemokines known as CCL17 and CCL22.

Fractalkine Human Recombinant produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 323 amino acids (25-339aa) and having a molecular mass of 34.3kDa. Fractalkine is fused to a 8 amino acid His tag at C-terminus and purified by proprietary chromatographic techniques.

Product Info

Amount : 2 µg / 10 µg

Purification : Greater than 90.0% as determined by SDS-PAGE.

Content : The Fractalkine solution (1 mg/ml) contains 10% Glycerol and Phosphate Buffered Saline (pH 7.4).

Storage condition : Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Amino Acid : QHHGVTKCNI TCSKMTSKIP VALLIHYQQN QASCGKRAII LETRQHRLFC ADPKEQWVKDAMQHLDRQAA
ALTRNGGTFE KQIGEVKPRTPAAGGMDES VVLEPEATGE SSSLEPTSSQEAQRALGTS PELPTGVTGS
SGTRLPPTPK AQDGGPVGTE LFRVPPVSTA ATWQSSAPHQPGPSLWAEAK TSEAPSTQDP STQASTASSP
APEENAPSEG QRVWGGQGSP RPENSLEREEMGPVPAHTDA FQDWGPGSMA HVSVPVSSE
GTPSREPVAS GSWTPKAEPP IHATMDPQRL GVLITVPVDA QAATRLEHHH HHH