

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-6777: GPT Rat

**Application:** Functional Assay

Alternative Name: Alanine aminotransferase 1 (EC:2.6.1.2), ALT1, Glutamate pyruvate transaminase 1, GPT 1, Glutamic-

-alanine transaminase 1, Glutamic--pyruvic transaminase 1, Gpt, Aat1, Gpt1.

## **Description**

Source: E.coli.

Sterile Filtered colorless solution.

GPT catalyzes the reversible transamination between alanine and 2-oxoglutarate to create pyruvate and glutamate. GPT has a crucial part in the intermediary metabolism of glucose and amino acids. GPT is broadly used as an indicator of liver reliability or hepatocellular destruction in clinical tests.

GPT Rat Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 519 amino acids (1-496 a.a) and having a molecular mass of 57.5kDa.Â

## **Product Info**

Amount:  $5 \mu g / 20 \mu g$ 

**Purification :** Greater than 90.0% as determined by SDS-PAGE.

**Content:** GPT protein solution (1mg/ml) containing Phosphate Buffered Saline pH 7.4 and 10% glycerol.

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods

**Storage condition:** of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Avoid multiple freeze-thaw cycles.

Amino Acid: MGSSHHHHHH SSGLVPRGSH MGSMASRVND QSQASRNGLK GKVLTLDTMN PCVRRVEYAV

RGPIVQRALE LEQELRQGVK KPFTEVIRAN IGDAQAMGQR PITFFRQVLA LCVYPNLLSS PDFPEDAKRR AERILQACGG HSLGAYSISS GIQPIREDVA QYIERRDGGI PADPNNIFLS TGASDAIVTM LKLLVSGEGR ARTGVLIPIP QYPLYSAALA ELDAVQVDYY LDEERAWALD IAELRRALCQ ARDRCCPRVL CVINPGNPTG QVQTRECIEA VIRFAFKEGL FLMADEVYQD NVYAEGSQFH SFKKVLMEMG PPYSTQQELA SFHSVSKGYM

GECGFRGGYV EVVNMDAEVQ KQMGKLMSVR LCPPVPGQAL MDMVVSPPTP SEPSFKQFQA

ERQEVLAELA AKAKLTEQVF NEAPGIRCNP VQGAMYSFPQ VQLPLKAVQR AQELGLAPDM FFCLCLLEET

GICVVPGSGF GOOEGTYHFR MTILPPMEKL RLLLEKLSHF HAKFTHEYS.

## **Application Note**

Specific activity is > 60units/mg, and is defined as the amount of enzyme that cleaves 1umole of L-Alanine to L-Glutamate per minute at pH 7.5 at 37C.