

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-6973: FGFR2 Human, (22-289)

EC 2.7.10, FGFR-2, BFR-1, CD332, BBDS, CEK3, ECT1, TK14, TK25, CFD1, KSAM, JWS, Fibroblast

Alternative Growth Factor Receptor 2, Keratinocyte Growth Factor Receptor, Bacteria-Expressed Kinase, EC 2.7.10.1, K-Name: SAM, KGFR, BEK, Protein Tyrosine Kinase, Receptor Like 14, BEK Fibroblast Growth Factor Receptor,

Craniofacial Dysostosis 1, Jackson-Weiss Syndrome, Pfeiffer Syndrome, Crouzon Syndrome, CD332 Antigen.

## **Description**

Source: Sf9, Baculovirus cells. Sterile Filtered colorless solution.

Fibroblast Growth Factors (FGFs) comprise a family of at least 18 structurally related proteins that take part in a multitude of physiological and pathological cellular processes, including differentiation, cell growth, angiogenesis, wound healing and tumorgenesis. The biological activities of the FGFs are mediated by a family if type I transmembrane tyrosine kinases which undergo dimerization and autophosphorylation after ligand binding. 4 different genes encoding related FGF receptors. Multiple forms of FGFR-1 to -3 are generated by alternative splicing of the mRNAs. A frequent splicing event involving FGFR-1 and -2 results in receptors containing all three Ig domains, referred to as the alpha isoform, or only IgII and IgIII. Only the alpha isoform has been identified for FGFR-3 and FGFR-4. Additional splicing events for FGFR-1 to -3, involving the C-terminal half of the IgIII domain encoded by 2 mutually exclusive alternative exons, generate FGF receptors with alternative IgIII domains (IIIb and IIIc). A IIIa isoform which is a secreted FGF binding protein containing only the N-terminal half of the IgIII domain plus some intron sequences has also been reported for FGFR-1.

FGFR2 produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 507 amino acids (22-289a.a.) and having a molecular mass of 56.8kDa. FGFR2 is expressed with a 239 amino acid hlgG-His-Tag at C-Terminus and purified by proprietary chromatographic techniques.

## **Product Info**

Amount:  $2 \mu g / 10 \mu g$ 

**Purification:** Greater than 95.0% as determined by SDS-PAGE.

Content: FGFR2 protein solution (0.5mg/ml) contains Phosphate Buffered Saline (pH 7.4) and 10%

glycerol

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of

Storage condition: time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid

multiple freeze-thaw cycles.

Amino Acid: RPSFSLVEDT TLEPEDAISS GDDEDDTDGA EDFVSENSNN KRAPYWTNTE

KMEKRLHAVPAANTVKFRCP AGGNPMPTMR WLKNGKEFKQ EHRIGGYKVR NQHWSLIMES VVPSDKGNYTCVVENEYGSI NHTYHLDVVE RSPHRPILQA GLPANASTVV GGDVEFVCKV YSDAQPHIQWIKHVEKNGSK YGPDGLPYLK VLKHSGINSS NAEVLALFNV TEADAGEYIC KVSNYIGQANQSAWLTVLPK QQAPGREKEI TASPDYLELE PKSCDKTHTC PPCPAPELLG GPSVFLFPPKPKDTLMISRT PEVTCVVVDV SHEDPEVKFN WYVDGVEVHN AKTKPREEQY NSTYRVVSVLTVLHQDWLNG KEYKCKVSNK ALPAPIEKTI SKAKGQPREP QVYTLPPSRD ELTKNQVSLTCLVKGFYPSD IAVEWESNGQ PENNYKTTPP VLDSDGSFFL YSKLTVDKSR

WQQGNVFSCSVMHEALHNHY TQKSLSLSPG KHHHHHH.