9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-13087: CD23 Human, Sf9

Alternative Name : Low affinity immunoglobulin epsilon Fc receptor isoform a, BLAST-2, CD23, CD23A, CLEC4J, FCE2, IGEBF, C-type lectin domain family 4 member J, Immunoglobulin E-binding factor, Lymphocyte IgE receptor.

Description

Source: Sf9, Baculovirus cells.
Sterile Filtered colorless solution.
FCER2, or Fc epsilon RII, is a receptor for IgE with low affinity, which is critical to regulation of IgE levels. IgE is an antibody isotype that attached to allergy and resistance to parasites. FCER2 is a C-type lectin, contrary to many of the receptors for antibodies. FCER2 is located on activated macrophages, mature B cells, follicular dendritic cells, platelets and eosinophils.
CD23Â produced in Sf9 Baculovirus cells is a single, glycosylated polypeptide chain containing 283 amino acids (48-321a.a.) and having a molecular mass of 32.0kDa.Â CD23Â is expressed with a 6 amino acid His tag at C-Terminus and purified by proprietary chromatographic techniques.

Product Info

Amount :	1 µg / 5 µg
Purification :	Greater than 90.0% as determined by SDS-PAGE.
Content :	CD23 protein solution (0.25mg/ml) contains phosphate buffered saline (pH7.4) and 10% glycerol.
Storage condition :	Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.
Amino Acid :	ADPDTTQSLK QLEERAARNV SQVSKNLESH HGDQMAQKSQ STQISQELEE LRAEQQRLKS QDLELSWNLN GLQADLSSFK SQELNERNEA SDLLERLREE VTKLRMELQV SSGFVCNTCP EKWINFQRKC YYFGKGTKQW VHARYACDDM EGQLVSIHSP EEQDFLTKHA SHTGSWIGLRNLDLKGEFIW VDGSHVDYSN WAPGEPTSRS QGEDCVMMRG SGRWNDAFCD RKLGAWVCDR LATCTPPASE GSAESMGPDS RPDPDGRLPTPSAPLHSHHH HHH.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.