

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-1684: Monoclonal Antibody to CD98 (SLC3A2)(Clone: UM7F8)

Clonality: Monoclonal **Clone Name:** UM7F8 FACS.IF Application: Reactivity: Human Gene: SLC3A2 Gene ID: 6520 **Uniprot ID:** P08195 Format: **Purified**

Alternative Name: SLC3A2, MDU1, 4F2 cell-surface antigen heavy chain, 4F2 heavy chain antigen, Lymphocyte

activation antigen 4F2 large subunit, Solute carrier family 3 member 2

Isotype: Mouse IgG1, kappa **Immunogen Information:** Molt 13 T cell line

Description

CD98 exits as a heterodimer containing a disulphide-linked glycosylated heavy chain and a non-glycosylated light chain. It is a member of the solute carrier family and encodes a cell surface, transmembrane protein. The protein exists as the heavy chain of a heterodimer, covalently bound through disulfide bonds to one of several possible light chains. The encoded transporter plays a role in regulation of intracellular calcium levels and transports L-type amino acids. Alternatively spliced transcript variants, encoding different isoforms, have been characterized.

Product Info

Amount: $100 \mu g$

Purification: Affinity Chromatography

Content: 100 μg in 500 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly

toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Comitogenic with soluble anti-CD2 and immobilized anti-CD3 mAbs; Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);