

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-13571: Borrelia Afzelii p100

Alternative Name : Borrelia belongs to a genus of bacteria of the spirochete phylum. Borrelia causes borreliosis, which is a zoonotic, vector-borne disease transmitted mainly by ticks and some by lice, depending on the species. Of the 36 known species of Borrelia, 12 are distinguished to cause Lyme disease or borreliosis and are transmitted by ticks. The main Borrelia species causing Lyme disease are Borrelia burgdorferi, Borrelia afzelii, and Borrelia garinii. The Borrelia genus members have a linear chromosome which is about 900 kbp in length as well as an excess of both linear and circular plasmids in the 5-220 kbp size range. The plasmids are atypical, as compared to most bacterial plasmids, since they contain many paralogous sequences, a large number of pseudogenes and, in some cases, essential genes. Moreover, a number of the plasmids have features suggesting that they are prophages.

Description

Source: Sf9 insect cells. Sterile Filtered clear solution.

Borrelia belongs to a genus of bacteria of the spirochete phylum. Borrelia causes borreliosis, which is a zoonotic, vector-borne disease transmitted mainly by ticks and some by lice, depending on the species. Of the 36 known species of Borrelia, 12 are distinguished to cause Lyme disease or borreliosis and are transmitted by ticks. The main Borrelia species causing Lyme disease are Borrelia burgdorferi, Borrelia afzelii, and Borrelia garinii. The Borrelia genus members have a linear chromosome which is about 900 kbp in length as well as an excess of both linear and circular plasmids in the 5-220 kbp size range. The plasmids are atypical, as compared to most bacterial plasmids, since they contain many paralogous sequences, a large number of pseudogenes and, in some cases, essential genes. Moreover, a number of the plasmids have features suggesting that they are prophages.

Recombinant Borrelia Afzelii Outer Surface Protein p100 produced in SF9 is a glycosylated, polypeptide chain having a calculated molecular mass of 74,782 Dalton. Borrelia Afzelii p100 is expressed with a 10xHis tag at N-terminus and purified by proprietary chromatographic techniques.

Product Info

Amount: 5 μg / 20 μg

Purification: Greater than 80.0% as determined by SDS-PAGE.

Content: Borrelia Afzelii p100 is supplied in 20mM HEPES buffer pH-7.6, 250mM NaCl and 20% glycerol.

Storage condition:

Storage condition:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods

of time. Avoid multiple freeze-thaw cycles.