

## 32-13605: HIV-1 p24

### Alternative Name :

Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus *Lentivirus*, part of the family of *Retroviridae*. *Lentiviruses* have many common morphologies and biological properties. Many species are infected by *lentiviruses*, which are characteristically responsible for long-duration illnesses with a long incubation period. *Lentiviruses* are transmitted as single-stranded, positive-sense, enveloped RNA viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.

### Description

Source: *Escherichia Coli*.

Sterile Filtered White lyophilized (freeze-dried) powder.

Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus *Lentivirus*, part of the family of *Retroviridae*. *Lentiviruses* have many common morphologies and biological properties. Many species are infected by *lentiviruses*, which are characteristically responsible for long-duration illnesses with a long incubation period. *Lentiviruses* are transmitted as single-stranded, positive-sense, enveloped RNA viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells. Recombinant HIV-1 p24 produced in *E. coli* is a non-glycosylated polypeptide chain having a molecular mass of 26.7kDa and fused to a His tag at C-terminus.

### Product Info

<b>Amount :</b>	100 µg / 0.5 mg
<b>Purification :</b>	Greater than 95.0% as determined by SDS-PAGE. Lyophilized from 1mg/ml in 20mM sodium carbonate pH-10.
<b>Content :</b>	It is recommended to reconstitute the lyophilized HIV-1 p24 in sterile 18M-cm H <sub>2</sub> O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.
<b>Storage condition :</b>	Store the lyophilized HIV-1 p24 between 2-8°C, do not freeze. Upon reconstitution HIV-1 p24 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.