

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-9025: Anti-SARS-CoV-2 RBD antibody(DM35), Rabbit mAb

Clone Name: DM35
Application: ELISA,FACS
Alternative Name: SARS-CoV-2 RBD
Isotype: Rabbit IgG

Immunogen Information: Recombinant SARS-CoV-2 (2019-nCoV) S protein RBD (Arg 319-Phe541) produced by using

human HEK293 cells

## **Description**

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.

## **Product Info**

Amount:  $100 \mu g$ 

**Purification:** Purified from cell culture supernatant by affinity chromatography

Preservative: 0.1% Procline 300

Content: Constituents: 50% Glycerol; PBS,pH 7.4; 0.1% BSA

Not Sterile

**Storage condition :** Store at -20°C for 12 months (Avoid repeated freezing and thawing)

## **Application Note**

Recommended Dilutions FACS 1/100




Figure 1. Elisa plate pre-coated by 2  $\mu$ g/ml (100 $\mu$ l/well) SARS-CoV-2 RBD protein can bind Rabbit Anti-SARS-CoV-2 RBD monoclonal antibody (clone:DM35) in a linear range of 0.19-200 ng/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2. Competition FACS assay demonstrating Rabbit anti-RBD monoclonal antibody (clone: DM35) blockade of SARS-CoV-2 (COVID-19) S protein RBD (1 $\mu$ g/ml, ) binding to Expi 293 cell line transfected with human ACE2. IC50=1013ng/ml. The Y-axis represents the geometric mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.