## **w** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

## 32-17121: Recombinant human CD40 Ligand Protein with N-Human Fc tag

Alternative Name : CD40LG,CD154,CD40L,HIGM1,IGM,IMD3,T-BAM,TNFSF5,TRAP,gp39

## Description

Expression Host : HEK293

The protein has a predicted molecular mass of 42.3 kDa after removal of the signal peptide. The apparent molecular mass of hFc-CD40 Ligand is approximately 40-53 kDa due to glycosylation.

The protein encoded by this gene is expressed on the surface of T cells. It regulates B cell function by engaging CD40 on the B cell surface. A defect in this gene results in an inability to undergo immunoglobulin class switch and is associated with hyper-IgM syndrome.

## **Product Info**

Amount :	50 µg
Purification :	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Content :	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Storage condition :	Store at -80°C for 12 months (Avoid repeated freezing and thawing)




Figure 1. Human CD40 Ligand Protein, hFc Tag on SDS-PAGE under reducing condition.

Figure 2. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human CD40 Ligand,hFc tagged protein can bind Human CD40, mFc-His tagged protein in a linear range of 0.51-320 ng/ml.