

## 32-17193: Recombinant human CXCR3 protein with C-terminal human Fc tag

**Alternative Name :** C-X-C chemokine receptor type 3, CXCR3, CXCR-3, CKR-L2?G protein-coupled receptor 9?CD183?IP-10 receptor

### Description

Expression Host : HEK293

The protein has a predicted molecular mass of 31.35 kDa after removal of the signal peptide.

This gene encodes a G protein-coupled receptor with selectivity for three chemokines, termed CXCL9/Mig (monokine induced by interferon-g), CXCL10/IP10 (interferon-g-inducible 10 kDa protein) and CXCL11/I-TAC (interferon-inducible T cell chemoattractant). Binding of chemokines to this protein induces cellular responses that are involved in leukocyte traffic, most notably integrin activation, cytoskeletal changes and chemotactic migration. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms (CXCR3-B) shows high affinity binding to chemokine, CXCL4/PF4 (PMID:12782716).

### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 50 µg   |
| <b>Purification :</b>      | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.              |
| <b>Content :</b>           | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization. |
| <b>Storage condition :</b> | Store at -80°C for 12 months (Avoid repeated freezing and thawing)  |