

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-17198: Recombinant human IL1B protein with C-terminal human Fc tag

Alternative Name: IL-1; IL1-BETA; IL1F2

## **Description**

Expression Host: HEK293

The protein has a predicted molecular mass of 42.35 kDa after removal of the signal peptide.

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2.

## **Product Info**

**Amount:** 50 μg

**Purification:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

**Content:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants

before lyophilization.

**Storage condition:** Store at -80°C for 12 months (Avoid repeated freezing and thawing)




Fig.1: Human IL1B Protein, hFc Tag on SDS-PAGE under reducing condition.




Fig.2: ELISA plate pre-coated by 1  $\hat{1}\frac{1}{4}$ g/mL (100  $\hat{1}\frac{1}{4}$ L/well) Human IL1B Protein, hFc Tag (32-17198) can bind Anti-IL1B Neutralizing antibody (12-2004) in a linear range of 0.61-156.25 ng/mL.