

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2505: Anti-MICA/MICB FITC (Clone: 6D4)

Clonality: Monoclonal

Clone Name : 6D4
Application : FACS
Reactivity : Human
Conjugate : FITC

Isotype: Mouse IgG2a

Immunogen Information: Transfected C1R cells expressing MICA

Description

MICA and MICB glycoproteins are members of MHC class I family, closely linked to HLA-B. However, unlike HLA molecules, MICA and MICB are not associated with beta2 microglobulin and are conformationally stable in the absence of conventional MHC class I peptide ligands. Both proteins are stress-induced antigens expressed mainly in gastrointestinal epithelium, where they are recognized by V-delta1 subset of gamma/delta T cells, and also on diverse epithelial tumor cells. Binding of MICA/MICB receptor, the NKG2D, leads to cytolytic response of NK cells, Tc cells, and gamma/delta T cells. Alternative splicing results in multiple isoforms, and some of them have been associated with susceptibility to psoriasis and psoriatic arthritis. Shedding of MICA-related antibodies and ligands is involved in the progression from monoclonal gammopathy of undetermined significance to multiple myeloma.

Specificity: The mouse monoclonal antibody 6D4 recognizes a common extracellular epitope on MICA and MICB glycoproteins, transmembrane ligands of NKG2D, and is able to block NKG2D-mediated activation of NK cells and cytotoxic T cells.

Product Info

Amount: 0.1 ma

Purification:

The purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum

conditions. The conjugate is purified by size-exclusion chromatography.

Content:

Formulation: Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide

Storage condition : Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1 : Separation of Jurkat cells stained using anti-human MICA/MICB (6D4) FITC antibody (concentration in sample 5 μg/ml, red-filled) from unstained Jurkat cells (black-dashed) in flow cytometry analysis (surface staining).