

### 30-2518: Anti-HLA-F Antibody (Clone : 3D11)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	3D11
<b>Application :</b>	FACS , WB
<b>Reactivity :</b>	Human
<b>Gene :</b>	HLA-F
<b>Gene ID :</b>	3134
<b>Format :</b>	Purified
<b>Alternative Name :</b>	CDA12,major histocompatibility complex, class I, F
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Inclusion body-derived HLA-F heavy chain

#### Description

HLA-F, an MHC class I molecule, is a type I transmembrane protein (heavy chain), which forms heterodimers with beta-2 microglobulin (light chain) and binds to KIR3DS1, KIR3DS4, KIR3DL2, ILT2, ILT4, and TAP. Unlike most other HLA proteins, HLA-F is mainly localized in the endoplasmic reticulum and Golgi apparatus, with only a small amount present on the cell surface in some cell types; surface expression can be induced by cell activation. It is thought to bind a restricted subset of peptides for immune presentation. Multiple transcript variants encoding different isoforms have been found for HLA-F gene. These variants lack a coding exon found in transcripts from other HLA paralogues due to an altered splice acceptor site, resulting in a shorter cytoplasmic domain.

**Specificity :** The mouse monoclonal antibody 3D11 recognizes an extracellular epitope of HLA-F, a 42 kDa type I transmembrane protein expressed on B cells, NK cells, monocytes, and T cells, but mainly in the endoplasmic reticulum and Golgi apparatus, only a small amount on the cell surface, where, however, it can be expressed after cell activation.

#### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography
<b>Content :</b>	1 mg/ml Formulation : Phosphate buffered saline (PBS) solution with 15 mM sodium azide
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.