

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2520: Anti-HLA-E Antibody (Clone: 3D12)

Clonality: Monoclonal

Clone Name: 3D12

Application: FACS, IP, ELISA

Reactivity: Human
Gene: HLA-E
Gene ID: 3133
Format: Purified

Alternative Name: HLA class I histocompatibility antigen, alpha chain E, MHC class I antigen E, major histocompatibility

complex, class I, E

Isotype: Mouse IgG1

Description

HLA-E (Human Leukocyte Antigen E) is a non-classical MHC I antigen, which is important for dialogue with NK cells and their regulation through interaction with CD94/NKG2 receptor. Like other MHC I molecules, transmembrane HLA-E molecule (45 kDa) associates with beta2 microglobulin. Unlike HLA-G, expression of HLA-E molecules is not so restricted, but it has been detected at least at mRNA level in virtually all cells and tissues examined. In peripheral blood, HLA-E protein is expressed at least in all mononuclear cells, but in different quantity (B cells and monocytes more than T cells and NK cells).

Specificity: The mouse monoclonal antibody 3D12 (also known as 3D12HLA-E) recognizes native extracellular part of HLA-E, an ubiquitously expressed non-classical MHC class I molecule, as well as free HLA-E.

Product Info

Amount: 0.1 mg

Purification: Purified from cell culture supernatant by protein-A affinity chromatography

Content: 1 mg/ml

Formulation: Phosphate buffered saline (PBS) solution with 15 mM sodium azide

Storage condition : Store at 2-8°C. Do not freeze.




Figure 1 : Flow cytometry surface staining pattern of human peripheral blood stained using anti-human HLA-E (3D12) purified antibody (concentration in sample 4 μ g/ml) GAM APC.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2 : Separation of human lymphocytes (red-filled) from blood debris (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human HLA-E (3D12) purified antibody (concentration in sample 4 μ g/ml) GAM APC.