

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2608: Anti-Human CD264 APC (Clone: TRAIL-R4-01)

Clonality: Monoclonal
Clone Name: TRAIL-R4-01

Application :FACSReactivity :HumanConjugate :APC

Gene : TNFRSF10D **Gene ID :** 8793

Alternative Name: TNFRSF10D, DcR2, TRUNDD, TRAILR4,TNF receptor superfamily member 10d

Isotype: Mouse IgG1

Immunogen Information: TRAIL-R4 (aa 1-210) - hlgGhc fusion protein

Description

TRAIL-R4 (CD264, TR4, DcR2, TRUNDD), expressed mainly on CD8+ and NK cells, belongs to receptors of TRAIL, a TNF-like membrane toxic protein that induces apoptosis in many tumour cells, but not in normal cells. TRAIL-R4, however, contains partially truncated death domain, thus it is unable to induce apoptosis and serves as a negative regulator of apoptotic signaling by impairment death-inducing signaling complex (DISC) processing. TRAIL-R4 interacts with death receptor 5 (DR5) in the native DISC in a TRAIL-dependent manner and prevents its corecruitment with death receptor 4 (DR4).

Specificity: The antibody TRAIL-R4-01 reacts with an extracellular epitope of TRAIL-R4, a 42 kDa transmembrane protein expressed on various blood cells.

Product Info

Amount: 0.1 mg

Purification: The purified antibody is conjugated with allophycocyanin (APC) under optimum conditions. The

conjugate is purified by size-exclusion chromatography.

0.1 mg/ml

Content : Formulation : Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium

azide

Storage condition: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1 : Flow cytometry surface staining pattern of CD264 transfected HEK-293 cells using anti-human CD264 (TRAIL-R4-01) APC (concentration in sample 1,67 μ g/ml).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2 : Separation of cells stained using anti-human CD264 (TRAIL-R4-01) APC antibody (concentration in sample 1,67 μ g/ml, red-filled) from CD264 cells stained using mouse IgG1 isotype control (MOPC-21) APC antibody (concentration in sample 1,67 μ g/ml, black-dashed) in flow cytometry analysis (surface staining) of CD264 transfected HEK-293 cell suspension.