

### 30-2646: Anti-Human CD158d Low Endotoxin Antibody (Clone : mAb#33)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	mAb#33
<b>Application :</b>	WB, ICC, Functional Assay, FACS , IP
<b>Reactivity :</b>	Human
<b>Gene :</b>	KIR2DL4
<b>Gene ID :</b>	3805
<b>Format :</b>	Low Endotoxin
<b>Alternative Name :</b>	KIR2DL4, KIR103AS, 103AS, 15.212, killer cell immunoglobulin like receptor, two Ig d
<b>Isotype :</b>	Mouse IgG1 kappa
<b>Immunogen Information :</b>	NK3.3 cells and KIR2DL4-Ig fusion protein

#### Description

CD158d / KIR2DL4 is a KIR family member that shares structural features with both activating and inhibitory receptors and may mediate different functions under different circumstances. It contains cytoplasmic ITIM, suggesting inhibitory function, but also transmembrane domain similar to those of activating KIRs. It has been reported that CD158d serves as an inhibitory receptor for peripheral and uterine NK cells, but its ligation with soluble mAbs (unlike immobilized mAbs) results in activation of IFN- $\gamma$  secretion. CD158d also binds both membrane form and soluble form of its ligand HLA-G.

**Specificity :** The mouse monoclonal antibody mAb#33 (also known as mAb 33 or 33) recognizes extracellular portion of CD158d / KIR2DL4, a 45 kDa NK cell marker. Cell surface expression and function of CD158d / KIR2DL4 depends on genotype of particular individuals.

#### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography
	1 mg/ml
<b>Content :</b>	Formulation : Low endotoxin azide free phosphate buffered saline (PBS) solution, 0.2 $\mu$ m filter sterilized. Endotoxin level is less than 0.01 EU/ $\mu$ g of the protein, as determined by the LAL test.
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.