

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2651: Anti-Human CD146 PerCP (Clone : P1H12)

Clone Name: Monoclonal P1H12
Application: FACS

Reactivity: Dog, Mouse, Human, Rabbit

Gene : MCAM
Gene ID : 4162

Alternative Name: MelCAM, MCAM, MUC18, melanoma cell adhesion molecule

Isotype: Mouse IgG1

Immunogen Information: cultured human umbilical cells

Description

CD146, also known as MCAM (melanoma cell adhesion molecule) or MUC18, is a heavily glycosylated transmembrane glycoprotein with more than 50% of the mass from carbohydrates. It is expressed on epithelial and endothelial cells, fibroblasts, multipotent mesenchymal stromal cells, activated T cells and activated keratinocytes, and on some cancer cells, especially melanoma. The presence of CD146 on circulating blood cells has been confined to the activated T cells rather than circulating endothelial cells. CD146 mediates heterophilic cell adhesion and regulates monocyte transendothelial migration.

Specificity: The mouse monoclonal antibody P1H12 recognizes an extracellular epitope of CD146, a 118 kDa transmembrane glycoprotein expressed on epithelial and endothelial cells, fibroblasts, multipotent mesenchymal stromal cells, melanoma cells, activated T cells and activated keratinocytes.

Product Info

Amount: 100 tests

Purification : The purified antibody is conjugated with PerCP under optimum conditions. The conjugate is

purified by size-exclusion chromatography.

Content: Formulation: Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium

azide

Storage condition: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 μ l reagent / 100 μ l of whole blood or 10 6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Separation of HUVEC cells stained using anti-human CD146 PerCP antibody from HUVEC cells stained using mouse IgG1 isotype control PerCP antibody in flow cytometry analysis of HUVEC cell suspension.