9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

30-2651: Anti-Human CD146 PerCP (Clone : P1H12)

Clonality :	Monoclonal
Clone Name :	P1H12
Application :	FACS
Reactivity :	Dog,Mouse,Human,Rabbit
Gene :	MCAM
Gene ID :	4162
Alternative Name :	MelCAM, MCAM, MUC18,melanoma cell adhesion molecule
Isotype :	Mouse IgG1
Immunogen Information :	cultured human umbilical cells

Description

CD146, also known as MCAM (melanoma cell adhesion molecule) or MUC18, is a heavily glycosylated transmembrane glycoprotein with more than 50% of the mass from carbohydrates. It is expressed on epithelial and endothelial cells, fibroblasts, multipotent mesenchymal stromal cells, activated T cells and activated keratinocytes, and on some cancer cells, especially melanoma. The presence of CD146 on circulating blood cells has been confined to the activated T cells rather than circulating endothelial cells. CD146 mediates heterophilic cell adhesion and regulates monocyte transendothelial migration.
Specificity : The mouse monoclonal antibody P1H12 recognizes an extracellular epitope of CD146, a 118 kDa transmembrane glycoprotein expressed on epithelial and endothelial cells, fibroblasts, multipotent mesenchymal stromal cells, melanoma cells, activated T cells and activated keratinocytes.

Product Info

Amount :	100 tests
Purification :	The purified antibody is conjugated with PerCP under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Content :	Formulation : Stabilizing phosphate buffered saline (PBS) solution containing 15 mM sodium azide
Storage condition :	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com




Figure 1: Separation of HUVEC cells stained using anti-human CD146 PerCP antibody from HUVEC cells stained using mouse IgG1 isotype control PerCP antibody in flow cytometry analysis of HUVEC cell suspension.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.