

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2662: Anti-Human CD109 Antibody (Clone: W7C5)

Clonality: Monoclonal
Clone Name: W7C5
Application: FACS
Reactivity: Human
Gene: CD109
Gene ID: 135228
Format: Purified

Alternative Name: CPAMD7, p180, r150, FLJ38569, GPA,CD109 molecule

Isotype: Mouse IgG1

Immunogen Information: WERI-RB-1 retinoblastoma cell line

Description

CD109, also known as the Gov platelet alloantigen, is a GPI-anchored glycoprotein which localizes to the surface of platelets, activated T-cells, and endothelial cells, as well as of various hematopoietic cells and T cell lines. The protein binds to and negatively regulates signaling by transforming growth factor beta (TGF-beta). Multiple transcript variants encoding different isoforms have been found for this gene. The Gov antigen system is involved in platelet transfusion reaction, posttransfusion purpura and in neonatal alloimmune thrombocytopenia.

Specificity: The mouse monoclonal antibody W7C5 recognizes CD109, an approximately 165 kDa GPI-anchored extracellular protein expressed mainly on various hematopoietic cells, activated T lymphoblasts, activated platelets, and endothelial cells.

Product Info

Amount: 0.1 mg

Purification : Purified from cell culture supernatant by protein-A affinity chromatography

Content:

Formulation: Phosphate buffered saline (PBS) solution with 15 mM sodium azide

Storage condition : Store at 2-8°C. Do not freeze.




Figure 1 : Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD109 (W7C5) purified antibody (concentration in sample 1 μ g/ml) GAM APC.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2 : Separation of human monocytes (red-filled) from CD109 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD109 (W7C5) purified antibody (concentration in sample 1 μ g/ml) GAM APC.