# **∗** abeomics

### 30-2664: Anti-Mouse CD106 Azide Free Antibody (Clone : 429 (MVCAM.A))

Clonality :	Monoclonal
Clone Name :	429 (MVCAM.A)
Application :	FACS , IP, IHC(F), Functional Assay
Reactivity :	Mouse
Gene :	Vcam1
Gene ID :	22329
Format :	Azide Free
Alternative Name :	Vcam-1,vascular cell adhesion molecule 1
Isotype :	Rat IgG2a kappa
Immunogen Information :	Murine preadipose cell line PA6

#### Description

CD106 / VCAM-1 (vascular cell adhesion molecule-1) is an Ig-like cell surface adhesion molecule binding VLA-4 integrin. VCAM-1 is a potent T cell costimulatory molecule taking part in their positive selection and survival, as well as in adhesion, transendothelial migration and activation of peripheral T cells. VCAM-1 is also involved in endothelial cell-cell contacts. Whereas VCAM-1 normally mediates leukocyte extravasion to sites of tissue inflammation, tumour cells can use overexpressed VCAM-1 to escape T cell immunity.

Specificity : The rat monoclonal antibody 429 (also known as MVCAM.A) recognizes an extracellular epitope of murine CD106, a 100-110 kDa type I membrane protein of the immunoglobulin superfamily, a crucial mediator of leukocyte adhesion, and a costimulation molecule.

#### **Product Info**

Amount : Purification :	0.1 mg Purified by protein-A affinity chromatography
Content :	1 mg/ml Formulation : Azide free phosphate buffered saline (PBS) solution, 0.2 $\mu$ m filter sterilized
Storage condition :	Store at 2-8°C. Do not freeze.



Figure 1 : Flow cytometry surface staining pattern of murine bone marrow cell suspension stained using anti-mouse CD160 (429) purified antibody (azide free, concentration in sample 0,19  $\mu$ g/ml) DAR APC.

# **₩** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure 2 : Separation of murine CD106 positive cells (red-filled) from CD106 negative cells (black-dashed) in flow cytometry analysis (surface staining) of murine bone marrow cell suspension stained using anti-mouse CD160 (429) purified antibody (azide free, concentration in sample 0,19  $\mu$ g/ml) DAR APC.