9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

12-8019: Anti-Human CD194 (CCR4) (Mogamulizumab)

Clonality :	Monoclonal
Clone Name :	KW-0761
Reactivity :	Human
Alternative Name :	CD194; CKR-4; CCR-4; CCR4; K5-5
Isotype :	Human IgG1k
Immunogen Information :	Humanization of mouse anti-CCR4 mAb7.

Description

Expression Host : HEK-293
Pathogen Testing : To protect mouse colonies from infection by pathogens and to assure that experimental preclinical data is not affected by such pathogens, all of this recombinant biosimilar antibodies are tested and guaranteed to be negative for all pathogens in the IDEXX IMPACT I Mouse Profile.
This non-therapeutic biosimilar antibody uses the same variable region sequence as the therapeutic antibody Mogamulizumab. Clone KW-0761 recognizes human CD194 (CCR4). This product is for research use only.
Clone KW-0761 (Mogamulizumab) is a research-grade, afucosylated, humanized monoclonal antibody generated from mouse anti-CCR4 mAb7 that targets human CCR4.1 CC chemokine receptor type 4 (CCR4) is a protein that belongs to the G protein-coupled receptor family and is a receptor for a variety of CC chemokines including MCP-1, MIP-1, RANTES, TARC, and Macrophage-derived chemokine. Chemokines are involved in the development, homeostasis, and function of the immune system and are known to regulate cell trafficking of various types of leukocytes. In a 2018 Phase I clinical trial, Mogamulizumab was found to decrease the number of HTLV-1Â–infected cells and the levels of inflammatory markers related to HTLV-1Â–Associated Myelopathy.3

Product Info

Amount :	50 µg / 100 µg
Purification :	>=95% monomer by analytical SEC
Content :	Concentration : >= 5.0 mg/ml This biosimilar antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added.
Storage condition :	Functional grade biosimilar antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at -80°C. Avoid Repeated Freeze Thaw Cycles.

Application Note

Endotoxin : <= 1.0 EU/mg as determined by the LAL method
The suggested concentration for Mogamulizumab biosimilar antibody for staining cells in flow cytometry is <= 0.25 µg per 106 cells in a volume of 100 µl. Titration of the reagent is recommended for optimal performance for each application.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.